arXiv:1208.2280v1 [math.RA] 10 Aug 2012

CONNECTED HOPF ALGEBRAS OF DIMENSION p^2

XINGTING WANG

Abstract. Let H be a finite-dimensional connected Hopf algebra and let p be a prime number. We provide the algebra structure of the associated graded Hopf algebra gr$H$. Then we consider the special case when H contains a normal Hopf subalgebra K and is generated by K and another element as an algebra. We also prove that H is local if and only if all the primitive elements in H are nilpotent, when H is cocommutative and the base field is $\mathbb{F}_p$. Finally we classify all connected Hopf algebras of dimension p^2 over an algebraically closed field.

1. Introduction

This paper serves as a beginning of classifying all connected Hopf algebras of dimension p^n for small integer $n \geq 1$ over an algebraically closed field $\mathbf{k}$. Let H be a finite-dimensional connected Hopf algebra as in [10, Def. 5.1.5]. In section 2, some basic definitions and properties are given concerning H. In section 3, we show the following algebra structure theorem, which applies especially to the associated graded Hopf algebra gr$H$.

Theorem 1.1 (Theorem 3.1). *Let* $H = \bigoplus_{n=0}^{\infty} H(n)$ *be a finite-dimensional connected coradically graded Hopf algebra. Then* H *is isomorphic to* $\mathbf{k}\left[x_1, x_2, \cdots, x_d\right] / \left(x_1^p, x_2^p, \cdots, x_d^p\right)$ *for some* $d \geq 0$ *as algebras.*

Section 4 treats the situation when H contains a normal Hopf subalgebra K [10, Def. 3.4.1] and is generated by K and another element as an algebra. The main result is highlighted in

Theorem 1.2 (Theorem 4.5). *Let* $K \subseteq H$ *be a degree* d *extension of finite-dimensional connected Hopf algebras. Suppose that* H *is generated by* K *and some* $x \in H$ *as an algebra. Also assume that* $\Delta(x) = x \otimes 1 + 1 \otimes x + u$ *where* $u \in K \otimes K$ *and* $[K, x] \subseteq \sum_{0 \leq i \leq 1} Kx^i$*. Then* H *is a free left* K*-module such that* $H = \bigoplus_{i=0}^{p^d-1} Kx^i$*. Furthermore if* K *is normal in* H*, then* x *satisfies a monic equation as follows:*

$$x^{p^d} + \sum_{i=0}^{d-1} a_i x^{p^i} + b = 0$$

for some $a_i \in \mathbf{k}$ *and* $b \in K$*.*

In section 5 we prove a special case of the conjecture that a finite-dimensional connected Hopf algebra is local if and only if all its primitive elements are nilpotent.

Theorem 1.3 (Theorem 5.4)**.** *Let* H *be a finite-dimensional cocommutative connected Hopf algebra over* $\mathbb{F}_p$*. Write* K *as the Hopf subalgebra generated by all primitive elements of* H*. Then the following are equivalent.*

1. H *is local.*
2. L *is local for some Hopf subalgebra* L *containing* K*.*
3. K *is local.*
4. *All the primitive elements in* H *are nilpotent.*

In the last section, we classify all connected Hopf algebras of dimension p^2 over an algebraically closed field.

Theorem 1.4 (Theorem 6.4)**.** *Let* H *be a connected Hopf algebra of dimension* p^2*. When* $\dim \mathrm{P}(H) = 2$*, it is isomorphic to one of the following:*

1. $\mathbf{k}\,[x,y]\,/\,(x^p, y^p)$*;*
2. $\mathbf{k}\,[x,y]\,/\,(x^p - x, y^p)$*;*
3. $\mathbf{k}\,[x,y]\,/\,(x^p - y, y^p)$*;*
4. $\mathbf{k}\,[x,y]\,/\,(x^p - x, y^p - y)$*;*
5. $\mathbf{k}\langle x,y\rangle/\,([xy] - y, x^p - x, y^p)$*,*

where x, y *are primitive. When* $\dim \mathrm{P}(H) = 1$*, it is isomorphic to one of the following:*

6. $\mathbf{k}\,[x,y]\,/(x^p, y^p)$*;*
7. $\mathbf{k}\,[x,y]\,/(x^p, y^p - x)$*;*
8. $\mathbf{k}\,[x,y]\,/(x^p - x, y^p - y)$*,*

where $\Delta\,(x) = x \otimes 1 + 1 \otimes x$ *and* $\Delta\,(y) = y \otimes 1 + 1 \otimes y + \sum_{i=1}^{p-1} \binom{p}{i}/p\; x^i \otimes x^{p-i}$*.*

Acknowledgment. The author thanks Professor James Zhang, Guangbin Zhuang and Cris Negron for their valuable suggestions. Also he appreciates the help from Professors Greenberg and Premet for some proofs in the paper. The research was partially supported by the US National Science Foundation.

2. Preliminaries

Throughout this paper, $\mathbf{k}$ denotes a base field, algebraically closed of characteristic $p \neq 0$. All vector spaces, algebras, coalgebras and tensor products are taken over $\mathbf{k}$ unless otherwise stated. Also V^* denotes the vector space dual of any vector space V. For any coalgebra C, the coradical C_0 is defined to be the sum of all simple subcoalgebras of C. It is called connected if C_0 is one-dimensional. Following [10, 5.2.1], $\{C_n\}_{n=0}^{\infty}$ is used to denote the coradical filtration of C. For any Hopf algebra H, we use $\mathrm{P}(H)$ to indicate the subspace of primitive elements. Following the terminology in [2, Def. 1.13], we recall the definition of graded Hopf algebras.

Definition 2.1. Let H be a Hopf algebra with antipode S. If

(1) $H = \bigoplus_{n=0}^{\infty} H(n)$ is a graded algebra,
(2) $H = \bigoplus_{n=0}^{\infty} H(n)$ is a graded coalgebra,
(3) $S(H(n)) \subseteq H(n)$ for any $n \geq 0$,

then H is called a graded Hopf algebra. If in addition,

(4) $H = \bigoplus_{n=0}^{\infty} H(n)$ is a coradically graded coalgebra,

then H is called a coradically graded Hopf algebra.

There are some basic properties of finite-dimensional Hopf algebras, which we use frequently.

Proposition 2.2. [9, 10] *Let H be a finite-dimensional Hopf algebra.*

(1) *H^* is a finite-dimensional Hopf algebra by dualizing all the maps.*
(2) *H is local if and only if H^* is connected.*
(3) *H is local, then any quotient or sub Hopf algebra of H is local.*

Furthermore assume that H is connected.

(4) *Any quotient or sub Hopf algebra of H is connected.*
(5) $\dim \mathrm{P}(H) = \dim J/J^2$*, where J is the Jacobson radical of H^*.*
(6) *H is primitively generated if and only if $H \cong u(\mathfrak{g})$, where $\mathfrak{g} = \mathrm{P}(H)$.*
(7) $\dim u(\mathfrak{g}) = p^{\dim \mathfrak{g}}$.
(8) $\dim H = p^n$ *for some integer n.*

Proof. (1) is from [10, Thm. 9.1.3].

(2), (5) are from [10, Prop. 5.2.9].

For (4) assume H is connected, H/I is connected by [10, Cor. 5.3.5], where I is any Hopf ideal of H. And for any Hopf subalgebra K of H, by [10, Lemma 5.2.12], $K_0 = K \bigcap H_0$. Since H_0 is one-dimensional, so is K_0. Thus K is connected.

(3) is the dual version of (4) by (1).

(6) is a standard result from [9] and (7) comes from [10, P. 23].

(8) is true because the associated graded ring $\mathrm{gr}_J(H^*)$ with respect to its J-adic filtration is connected and primitively generated. Hence $\dim H = \dim H^* = \dim \mathrm{gr}_J(H^*) = p^n$, where $n = \dim \mathrm{P}(\mathrm{gr}_J(H^*))$ by (7). □

Definition 2.3. Consider an extension of finite-dimensional connected Hopf algebras $K \subseteq H$.

1. If $\dim K = p^m$ and $\dim H = p^n$, then the degree of the extension is defined to be $n - m$.
2. The first order of the extension is defined to be the minimal integer n such that $K_n \subsetneq H_n$. And we say it is infinity if $K = H$.
3. The extension is said to be simple if H is generated by H_n as an algebra, where n is the first order of the extension.
4. The extension is said to be normal if K is a normal Hopf subalgebra of H.

Remark 2.4. By [10, Lemma 5.2.12] if D is a subcoalgebra of C, we have $D_n = D \bigcap C_n \subseteq C_n$. Also the coradical filtration is exhaustive for any coalgebra by [10, Theorem 5.2.2]. As a result of [10, Lemma 5.2.10], a connected bialgebra is automatically a connected Hopf algebra. Furthermore by [13, Lemma 6.1], we have that any sub-bialgebra of a connected Hopf algebra is a Hopf subalgebra. Let H be a connected Hopf algebra. Then the algebra generated by each coradical filtration H_n is a connected Hopf subalgebra of H. Because each coradical filtration H_n is a subcoalgebra and the algebra generated by it is certainly a sub-bialgebra.

Throughout the whole paper we would use the following convention:

Convention 2.5. Define the expression $\omega(x) = \sum_{i=1}^{p-1} \binom{p}{i}/p \; x^i \otimes x^{p-i}$, where $\binom{p}{i}/p \in \mathbf{k}$ for each $1 \le i \le p-1$.

3. Associated Graded Hopf algebras for finite-dimensional connected Hopf Algebras

Theorem 3.1. *Let $H = \bigoplus_{n=0}^{\infty} H(n)$ be a finite-dimensional connected coradically graded Hopf algebra. Then H is isomorphic to $\mathbf{k}\left[x_1, x_2, \cdots, x_d\right] / \left(x_1^p, x_2^p, \cdots, x_d^p\right)$ for some $d \ge 0$ as algebras.*

Proof. Denote $K = \bigoplus_{n=0}^{\infty} H(n)^*$ as the graded dual of H. It is a graded Hopf algebra and connected for $K_0 \subseteq K(0) = H(0)^* = \mathbf{k}$ by [10, Lemma 5.3.4]. Moreover since H is coradically graded, by [1, Lemma 5.5], K is generated in degree one and hence cocommutative. Therefore by duality H is commutative and local. Then according to [12, Theorem 14.4], H is isomorphic to $\mathbf{k}\,[x_1, x_2, \cdots, x_d]\,/(x_1^{p^{n_1}}, x_2^{p^{n_2}}, \cdots, x_d^{p^{n_d}})$ for some $d \geq 0$ as an algebra. Thus it suffices to prove inductively that for any homogeneous element $x \in H(n)$, we have $x^p = 0$ for all $n \geq 1$. Since H is coradically graded, $\mathrm{P}(H) = H(1)$. Then for any $x \in H(1)$, we have $x^p \in (H(1))^p \bigcap H(1) \subseteq H(p) \bigcap H(1) = 0$. Assume the assertion holds for $n \leq m-1$ and for any $x \in H(m)$. By the definition of graded Hopf algebras we have:

$$\Delta(x) = x \otimes 1 + 1 \otimes x + \sum_{i=1}^{m-1} y_i \otimes y_{m-i},$$

where $y_i \in H(i)$ for all $1 \leq i \leq m-1$. Therefore $\Delta(x^p) = x^p \otimes 1 + 1 \otimes x^p + \sum_{i=1}^{m-1} y_i^p \otimes y_{m-i}^p = x^p \otimes 1 + 1 \otimes x^p$ by induction. Thus $x^p \in (H(m))^p \bigcap H(1) \subseteq H(pm) \bigcap H(1) = 0$. □

Corollary 3.2. *The associated graded ring of a finite-dimensional connected Hopf algebra with respect to its coradical filtration is isomorphic to* $\mathbf{k}\,[x_1, x_2, \cdots, x_d]\,/\,(x_1^p, x_2^p, \cdots, x_d^p)$ *for some* $d \geq 0$ *as algebras.*

Proof. The associated graded space $\mathrm{gr}H = \bigoplus_{n \geq 0} H_n/H_{n-1}$ is a graded Hopf algebra by [10, P.62]. Also mentioned in [2, Def. 1.13], $\mathrm{gr}H$ is coradically graded. Therefore $\mathrm{gr}H$ is a coradically graded Hopf algebra, which is clearly connected because H is connected. Hence $\mathrm{gr}H$ satisfies all the conditions in Theorem 3.1 and the result follows. □

As a consequence of the commutativity of the associated graded ring for any finite-dimensional connected Hopf algebra we conclude that:

Corollary 3.3. *Let H be a finite-dimensional connected Hopf algebra. Then* $[H_n, H_m] \subseteq H_{n+m-1}$ *for all integers n, m.*

4. Extensions of finite-dimensional Connected Hopf algebras

Lemma 4.1. *Let $K \subseteq H$ be an extension of finite-dimensional connected Hopf algebras with first order n. Then the degree of the extension is greater or equal to* $\dim(H_n/K_n)$.

Proof. By [10, Lemma 5.2.12], the inclusion $K \hookrightarrow H$ induces an injection $K_i/K_{i-1} \hookrightarrow H_i/H_{i-1}$ for all $i \geq 1$. Thus $\mathrm{gr}K = \bigoplus_{i\geq 0} K(i) \hookrightarrow \mathrm{gr}H = \bigoplus_{i\geq 0} H(i)$ and $K(i) = H(i)$ for all $0 \leq i \leq n-1$ since n is the first order of the extension. Moreover by [2, Def. 1.13], $(\mathrm{gr}H)_m = \bigoplus_{0\leq i\leq m} H(m)$ for all $m \geq 0$ and the same is true for $\mathrm{gr}K$. Therefore it is enough to prove the result in the associated graded Hopf algebras extension $\mathrm{gr}K \subseteq \mathrm{gr}H$.

For simplicity, we write K for $\mathrm{gr}K$, H for $\mathrm{gr}H$ and use $\mathrm{d}(H/K)$ to denote the degree of the extension $K \subseteq H$. We will prove the result by induction on $\dim(H_n/K_n)$. When $\dim(H_n/K_n) = 1$, it is trivial. Now suppose that $\dim(H_n/K_n) > 1$ and choose any $x \in H(n) \setminus K(n)$. Because H is a graded coalgebra,

$$\Delta(x) = x \otimes 1 + 1 \otimes x + \sum_{i=1}^{n-1} y_i \otimes y_{n-i},$$

where $y_i \in H(i) = K(i)$ for all $1 \leq i \leq n-1$. Hence K and x generate a Hopf subalgebra of H by Remark 2.4, which we denote as L. Now according to Corollary 3.2, we have $x^p = 0$. Thus $K \subseteq L$ is a degree one extension with first order n. Because H is a graded algebra, it is clear that L_n is spanned by K_n and x. Hence $\dim(L_n/K_n) = 1$ and $\dim(H_n/L_n) = \dim(H_n/K_n) - 1$. Therefore by induction we have $\dim(H_n/K_n) = \dim(H_n/L_n) + \dim(L_n/K_n) = \dim(H_n/L_n) + 1 \leq \mathrm{d}(H/L) + 1 = \mathrm{d}(H/L) + \mathrm{d}(L/K) = \mathrm{d}(H/K)$. □

Lemma 4.2. *Let $K \subseteq H$ be a simple extension with first order n. Then K is normal in H if and only if $[K, H_n] \subseteq K$.*

Proof. First suppose that K is normal in H. By [10, Lemma 5.3.2] for any $x \in H_n$, $\Delta(x) - x \otimes 1 - 1 \otimes x \in H_{n-1} \otimes H_{n-1} = K_{n-1} \otimes K_{n-1} \subseteq K \otimes K$. Thus we can write $\Delta(x) = x \otimes 1 + 1 \otimes x + \sum a_i \otimes b_i$ where $a_i, b_i \in K$. Apply the antipode S to get

$$S(x) = \varepsilon(x) - x - \sum a_i S(b_i).$$

By the definition of normal Hopf subalgebras [10, Def. 3.4.1], for any $y \in K$

$$\sum x_1 y S(x_2) = xy + yS(x) + \sum a_i y S(b_i) = u \in K.$$

Therefore

$$[y, x] = yx - xy = y\left(\varepsilon(x) - \sum a_i S(b_i)\right) + \sum a_i y S(b_i) - u \subseteq K,$$

which shows that $[K, H_n] \subseteq K$. Conversely suppose that $[K, H_n] \subseteq K$. Then it is clear that $K^+ H_n \subseteq H_n K^+ + K^+ \subseteq HK^+$ since $[K^+, H_n] \subseteq K^+$. We claim that $K^+(H_n)^i \subseteq HK^+$ for all

$i \geq 0$ by induction. Suppose the inclusion holds for i and then for $i+1$:

$$K^+ (H_n)^{i+1} = K^+ (H_n)^i H_n \subseteq \left(HK^+\right) H_n \subseteq H \left(HK^+\right) \subseteq HK^+.$$

Therefore $K^+H = \bigcup K^+(H_n)^i \subseteq HK^+$ and by symmetry $K^+H = HK^+$. According to [10, Cor. 3.4.4], K is normal. $\square$

Lemma 4.3. *Let H be a Hopf algebra with subalgebra K. If $x \in H$ satisfies $[K,x] \subseteq K$ and $\Delta(x) - x\otimes 1 - 1 \otimes x \in K \otimes K$, then $\Delta\left(x^{p^n}\right) - x^{p^n} \otimes 1 - 1 \otimes x^{p^n} \in K \otimes K$ for all $n \geq 0$.*

Proof. Denote $\Delta(x) = x\otimes 1 + 1\otimes x + u$ where $u \in K\otimes K$. First we prove that $\Delta(x^p) - x^p\otimes 1 - 1\otimes x^p \in K \otimes K$. By Lemma A.1 we have:

$$\Delta\left(x^p\right) = (x\otimes 1 + 1\otimes x + u)^p = x^p \otimes 1 + 1 \otimes x^p + u^p + \sum_{i=1}^{p-1} S_i$$

where iS_i is the coefficient of λ^{i-1} in $u\left(\operatorname{ad}\left(\lambda u + x\otimes 1 + 1 \otimes x\right)\right)^{p-1}$. Hence it suffices to show inductively that

$$u\left(\operatorname{ad}\left(\lambda u + x\otimes 1 + 1\otimes x\right)\right)^n \in (K\otimes K)[\lambda]$$

for all $n \geq 0$. Notice that when $n = 0$, it is just the assumption. Suppose it's true for $n-1$ then for n

$$\begin{aligned} u\left(\operatorname{ad}\left(\lambda u + x\otimes 1 + 1\otimes x\right)\right)^n &\in [(K\otimes K)[\lambda], \lambda u + x\otimes 1 + 1\otimes x] \\ &\subseteq \{[K\otimes K, u] + [K,x]\otimes K + K \otimes [K,x]\}[\lambda] \\ &\subseteq (K\otimes K)[\lambda]. \end{aligned}$$

Now replace x with $x^{p^{n-1}}$ and we have $[K, x^{p^{n-1}}] = K\left(\operatorname{ad}(x)\right)^{p^{n-1}} \subseteq K$ by Lemma A.1. The other cases can be proved in the similar way. $\square$

Let K be a Hopf subalgebra of H. Suppose there exists some $x \in H$ such that $\Delta(x) = x\otimes 1 + 1\otimes x + u$ where $u \in K\otimes K$ and $[K,x] \subseteq \sum_{0\leq i\leq 1} Kx^i$. Denote $L_n = \sum_{i\leq n} Kx^i$ for all $n \geq 0$. Then we obtain the following lemma:

Lemma 4.4. *L_n is a subcoalgebra and K-bimodule for all $n \geq 0$. In particular if H is generated by K and x as an algebra, then $H = \bigcup_{n\geq 0} L_n$.*

Proof. We first prove that $[K, x^n] \subseteq \sum_{0 \le i \le n} Kx^i$ for all $n \geq 0$ by induction on n. Certainly by assumption it holds for $n = 1$. Suppose it is true for $n - 1$. Therefore for n

$$\begin{aligned}
[a, x^n] &= (ax^{n-1})x - x^n a \\
&\in \left(x^{n-1}a + \sum_{0 \le i \le n-1} Kx^i \right) x - x^n a \\
&\subseteq x^{n-1} \left(xa + \sum_{0 \le i \le 1} Kx^i \right) + \sum_{1 \le i \le n} Kx^i - x^n a \\
&\subseteq \sum_{0 \le i \le 1} \left(Kx^{n-1} + \sum_{0 \le j \le n-1} Kx^j \right) x^i + \sum_{1 \le i \le n} Kx^i \\
&\subseteq \sum_{0 \le i \le n} Kx^i
\end{aligned}$$

for any $a \in K$. Thus $[K, x^n] \subseteq \sum_{0 \le i \le n} Kx^i$ for all $n \geq 0$.

It is clear that $L_n = \sum_{0 \le i \le n} Kx^i$ is a left K-module. The above result ensures that $L_n K \subseteq L_n$. Hence L_n is a K-bimodule for all $n \geq 0$. To prove L_n is a subcoalgebra, again by induction we first show that $\Delta(x^n) - x^n \otimes 1 - 1 \otimes x^n \in L_{n-1} \otimes L_{n-1}$ for all $n \geq 1$. When $n = 1$, it is just the assumption. Suppose it's true for $n - 1$ and write $\Delta(x^{n-1}) = x^{n-1} \otimes 1 + 1 \otimes x^{n-1} + \sum a_i \otimes b_i$ where $\sum a_i \otimes b_i \in L_{n-2} \otimes L_{n-2}$. Therefore

$$\begin{aligned}
& \Delta(x^n) - x^n \otimes 1 - 1 \otimes x^n \\
= \ & (x \otimes 1 + 1 \otimes x + u) \left(x^{n-1} \otimes 1 + 1 \otimes x^{n-1} + \sum a_i \otimes b_i \right) - x^n \otimes 1 - 1 \otimes x^n \\
\in \ & x \otimes x^{n-1} + x^{n-1} \otimes x + xL_{n-2} \otimes L_{n-2} + L_{n-2} \otimes xL_{n-2} + L_{n-2} \otimes L_{n-2} \\
\subseteq \ & L_{n-1} \otimes L_{n-1}.
\end{aligned}$$

Then it is straightforward to check that L_n is a coalgebra:

$$\begin{aligned}
\Delta(L_n) &\subseteq \Delta\left(\sum_{0\le i\le n} Kx^i\right)\\
&\subseteq \sum_{0\le i\le n} \Delta(K)\Delta\left(x^i\right)\\
&\subseteq \sum_{0\le i\le n} (K\otimes K)\left(x^i\otimes 1 + 1\otimes x^i + L_{i-1}\otimes L_{i-1}\right)\\
&\subseteq L_n\otimes L_n.
\end{aligned}$$

Furthermore if H is generated by K and x as an algebra, it is easy to see $H=\bigcup_{n\ge 0} L_n$. □

Theorem 4.5. *Let $K\subseteq H$ be a degree d extension of finite-dimensional connected Hopf algebras. Suppose that H is generated by K and some $x\in H$ as an algebra. Also assume that $\Delta(x)=x\otimes 1+1\otimes x+u$ where $u\in K\otimes K$ and $[K,x]\subseteq \sum_{0\le i\le 1} Kx^i$. Then H is a free left K-module such that $H=\bigoplus_{i=0}^{p^d-1} Kx^i$. Furthermore if K is normal in H, then x satisfies a monic equation as follows:*

$$x^{p^d}+\sum_{i=0}^{d-1} a_i x^{p^i}+b=0$$

for some $a_i\in \mathbf{k}$ and $b\in K$.

Proof. Denote $L_n=\sum_{0\le i\le n} Kx^i$ for all $n\ge 0$. By the previous Lemma 4.4, L_n is a subcoalgebra. Also H is a left K-module with generators $\{x^i|i\ge 0\}$ for $H=\sum Kx^i$. Because H is finite-dimensional, there exist some nontrivial relations between the generators such as

$$d_m x^m + d_{m-1}x^{m-1}+\cdots+d_1x+d_0=0,$$

where $d_i\in K$ and $d_m\neq 0$, among which we choose the lowest degree in terms of x, say degree m. Furthermore denote $D=K$, $L=L_{m-1}$, $F=x^m$ and $V=\{a\in D|aF\in L\}$. As in the proof of Lemma 4.4, we know $\Delta(F)-x^m\otimes 1-1\otimes x^m\in L\otimes L$. Then D,L,F satisfy all the conditions listed in [11, Lemma 1.1]. Hence $V=D$ for $0\neq d_m\in V$. Thus $x^m\in\bigoplus_{i<m} Kx^i$ and consequently H is a free left K-module with the free basis $\{x^i|0\le i\le m-1\}$. Since $\dim H=m\dim K$, it is easy to see $m=p^d$ by definition.

Now assume that K is normal. Follow the proof in Lemma 4.2, we can show that $[K, x] \subseteq K$. From pervious discussion there exists a general equation for x:

$$x^{p^d} + \sum_{i=0}^{p^d-1} a_i x^i = 0, \tag{4.0.1}$$

where all $a_i \in K$. According to Lemma 4.3, we can write $\Delta\left(x^{p^n}\right) = x^{p^n} \otimes 1 + 1 \otimes x^{p^n} + u_n$ where $u_n \in K \otimes K$ for all $n \geq 0$. Now apply the comultiplication Δ to the above identity (4.0.1) to get

$$x^{p^d} \otimes 1 + 1 \otimes x^{p^d} + u_d + \sum_{i=0}^{p^d-1} \Delta(a_i)(x \otimes 1 + 1 \otimes x + u)^i = 0.$$

Replacing x^{p^d} with $\left(-\sum_{i=0}^{p^d-1} a_i x^i\right)$, the following equation is straightforward:

$$\begin{aligned}\left(-\sum_{i=0}^{p^d-1} a_i x^i\right) \otimes 1 + 1 \otimes \left(-\sum_{i=0}^{p^d-1} a_i x^i\right) + \sum_{i=0}^{d-1} \Delta\left(a_{p^i}\right)\left(x^{p^i} \otimes 1 + 1 \otimes x^{p^i} + u_i\right) \\ + \sum_{i \in S} \Delta\left(a_i\right)(x \otimes 1 + 1 \otimes x + u)^i + \Delta(a_0) + u_d = 0\end{aligned} \tag{4.0.2}$$

with the index set $S = \{1, 2, \cdots, p^d\} \setminus \{1, p, p^2, \cdots, p^d\}$.

We first prove that $a_i = 0$ for all $i \in S$ by contradiction. If not suppose $n \in S$ is the largest integer such that $a_n \neq 0$. The free K-module structure for H implies that the $K \otimes K$-module $H \otimes H$ has a free basis $\left\{x^i \otimes x^j | 0 \leq i, j < p^d\right\}$. Thus the term $Kx^{n-i} \otimes Kx^i$ would only come from $\Delta\left(a_n\right)(x \otimes 1 + 1 \otimes x + u)^n$ for all $1 \leq i \leq n-1$. Moreover it exactly comes from $\Delta\left(a_n\right)(x \otimes 1 + 1 \otimes x)^n$ by the choice of n. Therefore $\binom{n}{i}\Delta\left(a_n\right)\left(x^{n-i} \otimes x^i\right) = 0$ for all $1 \leq i \leq n-1$. Suppose $n = p^\alpha m$ where $m > 1$ and $m \not\equiv 0 \pmod{p}$. Choose $i = p^\alpha$. Hence by [5, Lemma 5.1], $\binom{n}{p^\alpha} \equiv m \pmod{p}$. Then $\Delta(a_n) = 0$, which implies that $a_n = 0$, a contradiction. Therefore from equation (4.0.2), we deduce that $\Delta(a_{p^i})(x^{p^i} \otimes 1) = a_{p^i} x^{p^i} \otimes 1$ for all $0 \leq i \leq d-1$. Thus $\Delta(a_{p^i}) = a_{p^i} \otimes 1$. Then since H is counital, all of a_{p^i} are coefficients in the base field $\mathbf{k}$. □

5. FINITE-DIMENSIONAL COCOMMUTATIVE LOCAL CONNECTED HOPF ALGEBRAS OVER $\mathbb{F}_p$

Throughout this section we assume that $\mathbf{k} = \mathbb{F}_p$.

Lemma 5.1. *Let H be a finite-dimensional Hopf algebra with normal Hopf subalgebras $K \subseteq L \subseteq H$. Then there exists a natural isomorphism:*

$$\left(H/K^+H\right)^* \Big/ \left(H/L^+H\right)^{*+} \left(H/K^+H\right)^* \cong \left(L/K^+L\right)^*.$$

Proof. By [10, Theorem 2.1.3], L is Frobenius. Hence the injective left L-module map $L \hookrightarrow H$ splits since L is self-injective. Therefore we can write $H = L \bigoplus M$ as a direct sum of two left L-modules. Because $K \subseteq L$, we have $L \bigcap K^+H = L \bigcap K^+ (L \bigoplus M) = L \bigcap (K^+L \bigoplus K^+M) = K^+L$. Then the inclusion map $L \hookrightarrow H$ induces an injective Hopf algebra map $L/K^+L \hookrightarrow H/K^+H$, since K^+L and K^+H are Hopf ideals of L and H by [10, Lemma 3.4.2].

It is clear that the composition map $L/K^+L \hookrightarrow H/K^+L \twoheadrightarrow H/L^+H$ factors through $\mathbf{k}$ by the counit. Thus the dualized map restricted on $(H/L^+H)^{*+} = (H/L^+H)^* \bigcap \mathrm{Ker} u^* \to (L/K^+L)^*$ is the zero map where u is the unit map in H.

Therefore the natural surjective map $(H/K^+H)^* \twoheadrightarrow (L/K^+L)^*$, which is induced by the inclusion $L/K^+L \hookrightarrow H/K^+H$, factors through $\left(H/K^+H\right)^* \Big/ \left(H/L^+H\right)^{*+} \left(H/K^+H\right)^*$. In order to show that it is an isomorphism, it is enough to prove that both sides have the same dimension. By [10, Theorem 3.3.1], we have

$$\begin{aligned}
\dim \left(H/K^+H\right)^* \Big/ \left(H/L^+H\right)^{*+} \left(H/K^+H\right)^* &= \dim \left(H/K^+H\right)^* \Big/ \dim(H/L^+H)^* \\
&= (\dim H/\dim K) \Big/ (\dim H/\dim L) \\
&= \dim L/\dim K \\
&= \dim(L/K^+L)^*.
\end{aligned}$$

□

Proposition 5.2. *Let H be a finite-dimensional cocommutative connected Hopf algebra. Then H has a sequence of increasing normal Hopf subalgebras: $\mathbf{k} = K_0 \subset K_1 \subset \cdots \subset K_n = H$ satisfying the following properties:*

1. *Denote J as the Jacobson radical of H^*. Then the length n is the minimal integer such that $x^{p^n} = 0$ for all $x \in J$.*
2. *K_1 is the Hopf subalgebra of H generated by all primitive elements.*
3. *There are natural injective Hopf algebra maps:*

$$K_n/K_{n-1}^+K_n \hookrightarrow K_{n-1}/K_{n-2}^+K_{n-1} \hookrightarrow \cdots \hookrightarrow K_1/K_0^+K_1.$$

4. $0 = \dim \mathrm{P}\left(H/K_n^+H\right) \leq \dim \mathrm{P}\left(H/K_{n-1}^+H\right) \leq \cdots \leq \dim \mathrm{P}\left(H/K_0^+H\right) = \dim \mathrm{P}(H)$.

Proof. (1) By duality H^* is a finite-dimensional commutative local Hopf algebra. Therefore by [12, Theorem 14.4] we can write:

$$H^* = \mathbf{k}\left[x_1, x_2, \cdots, x_d\right] \Big/ \left(x_1^{p^{n_1}}, x_2^{p^{n_2}}, \cdots, x_d^{p^{n_d}}\right)$$

for some $d \geq 0$, in which we can define a decreasing chain of normal Hopf ideals [10, Def. 3.4.5]

$$\left(J_m = (x_1^{p^m}, x_2^{p^m}, \cdots, x_d^{p^m})\right)_{m\geq 0}.$$

By [10, P. 36], in the dual vector space H we have an increasing sequence of normal Hopf subalgebras: $\mathbf{k} = K_0 \subset K_1 \subset \cdots \subset K_m \subseteq \cdots \subseteq H$, where $K_m = (H^*/J_m)^*$ for all $m \geq 0$. For the length of this filtration, notice that $K_m = H \Leftrightarrow J_m = 0 \Leftrightarrow x_i^{p^m} = 0$ for all $1 \leq i \leq d \Leftrightarrow x^{p^m} = 0$ for all $x \in J_0 = J$.

(2) Write L as the Hopf subalgebra of H generated by $\mathrm{P}(H)$. By [10, Prop. 5.2.9], $\mathbf{k} \bigoplus \mathrm{P}(H) = \{h \in H | \langle J^2, h\rangle = 0\}$. Hence under the natural identification, $\mathrm{P}(H) \subset (H^*/J^2)^* \subseteq (H^*/J_1)^* = K_1$. Because L is generated by $\mathrm{P}(H)$ as an algebra, we have $L \subseteq K_1$. Moreover we know $\dim L = p^{\dim \mathrm{P}(H)} = p^{\dim J/J^2} = p^d$. On the other side $\dim K_1 = \dim H^*/J_1 = p^d$, which implies that $L = K_1$.

(3) Define a decreasing series of normal Hopf algebras of H^* by

$$A_m = \{h^{p^m} | h \in H^*\} = \mathbf{k}\left[x_1^{p^m}, x_2^{p^m}, \cdots, x_d^{p^m}\right].$$

Notice that $A_m^+ H^* = J_m$ for all $m \geq 0$. By Lemma 5.1 we have

$$\begin{aligned}\left(A_m/A_{m+1}^+ A_m\right)^* &\cong \left(H^*/A_{m+1}^+ H^*\right)^* \Big/ \left(H^*/A_m^+ H\right)^{*+} \left(H^*/A_{m+1}^+ H^*\right)^* \\ &= K_{m+1} \Big/ {K_m}^+ K_{m+1}.\end{aligned}$$

It is easy to see there exists a series of surjective p-power Hopf algebra maps $\phi_m : A_m \twoheadrightarrow A_{m+1}$ such that $\phi_m(x) = x^p$ for all $x \in A_m$. And it induces a series of surjective maps on the quotients $A_m/A_{m+1}^+ A_m \twoheadrightarrow A_{m+1}/A_{m+2}^+ A_{m+1}$ for all $m \geq 0$. By daulizing all the maps and the above natural isomorphism we have a series of injective maps

$$K_n/K_{n-1}^+ K_n \hookrightarrow K_{n-1}/K_{n-2}^+ K_{n-1} \hookrightarrow \cdots \hookrightarrow K_1/K_0^+ K_1.$$

(4) In Lemma 5.1, let $K = \mathbf{k}$ and $L = A_m$ then we have the special isomorphism:

$$A_m^* \cong H \Big/ K_m^+ H.$$

Therefore by Proposition 2.2(5),

$$\dim \mathrm{P}(H/K_m^+H) = \dim J(A_m)/J(A_m)^2 = \#\left\{\{x_1^{p^m}, x_2^{p^m}, \cdots, x_d^{p^m}\} \setminus \{0\}\right\},$$

which is the number of generators among $\{x_1, x_2, \cdots, x_d\}$ whose p^m-th power does not vanish. Thus the inequalities follow. □

Remark 5.3. When H is a finite-dimensional cocommutative connected Hopf algebra over any field of characteristic p, then all the results in Proposition 5.2 remain true except for (3), where we would only have injective coalgebra maps. In particular suppose that $\dim \mathrm{P}(H) = 1$ and $\dim H = p^n$. It is clear that $H^* \cong \mathbf{k}[x] / \left(x^{p^n}\right)$ as and algebra and has Jacobson radical $J = (x)$. Hence H is a finite-dimensional cocommutative connected Hopf algebra. Hence it can be filtered by an increasing sequence of normal Hopf subalgebras: $\mathbf{k} = K_0 \subset K_1 \subset \cdots \subset K_n = H$ such that K_1 is generated by primitive elements and $\dim K_m = p^m$ for all $0 \le m \le n$.

Theorem 5.4. *Let H be a finite-dimensional cocommutative connected Hopf algebra over $\mathbb{F}_p$. Write K as the Hopf subalgebra generated by all primitive elements of H. Then the following are equivalent.*

1. *H is local.*
2. *L is local for some Hopf subalgebra L containing K.*
3. *K is local.*
4. *All the primitive elements in H are nilpotent.*

Proof. (1) ⇒ (2) is from Proposition 2.2(3) and (2) ⇒ (3) is trivial. (3) ⇒ (4) is clear since K contains all the primitive elements in H and the augmentation ideal of K is nilpotent.

In order to show that (4) ⇒ (3), denote $\mathfrak{g} = \mathrm{P}(H)$ which is a restricted Lie algebra. Then (4) is equivalent to the statement that $\mathfrak{g}^{p^n} = 0$ for sufficient larger n. Therefore $(\mathrm{ad}x)^{p^n} = \mathrm{ad}(x^{p^n}) = 0$ for all $x \in \mathfrak{g}$. By Engel's Theorem [4, §3.2], g is nilpotent. Any representation of $K = u(\mathfrak{g})$ is a restricted representation of $\mathfrak{g}$. Therefore any irreducible representation of K is one-dimensional with trivial action of the augmentation ideal of K. Hence the augmentation ideal of K is nilpotent and K is local.

Finally we only need to show (3) ⇒ (1). First H is local if and only if the Jacobson radical of H equals $\ker \varepsilon$, where ε is the counit of H. We do induction on the dimension of H. When $\dim H = p$, the result is clear since $H = K$. By Proposition 5.2(3), there exists some proper normal Hopf subalgebra $L \supseteq K$ such that $H/L^+H \hookrightarrow K$ as Hopf subalgebra. Hence by Proposition 2.2(3) again, H/L^+H is local. Therefore there exists some integer l such that $(\ker \varepsilon)^l \subseteq L^+H$. By induction L is local. Then there is another integer s such that $(L^+)^s = 0$. Thus $(\ker \varepsilon)^{ls} \subseteq (L^+H)^s = (L^+)^s H = 0$. Here we use the fact that $L^+H = HL^+$ in the case that L is normal by [10, Cor. 3.4.4]. Since the Jacobson radical contains any nilpotent ideal, $J(H) = \ker \varepsilon$ and H is local. $\square$

6. Connected Hopf algebras of dimension p^2

The starting point for classifying finite-dimensional connected Hopf algebras turns out to be when the dimension is just p, where it is obvious that such Hopf algebras are primitively generated, i.e., by some primitive element x. As a consequence of the characteristic of the base field, x^p is still primitive which implies that $x^p = \lambda x$ for some $\lambda \in \mathbf{k}$ since the dimension of the primitive space is one. By rescaling of the variable we can always assume the coefficient λ is zero or one. Thus we have the following result:

Theorem 6.1. *All connected Hopf algebras of dimension p are isomorphic to either $\mathbf{k}[x]/(x^p)$ or $\mathbf{k}[x]/(x^p - x)$ where x is primitive.*

Corollary 6.2. *All local Hopf algebras of dimension p are isomorphic to $\mathbf{k}[x]/(x^p)$ with comultiplication either $\Delta(x) = x \otimes 1 + 1 \otimes x$ or $\Delta(x) = x \otimes 1 + 1 \otimes x + x \otimes x$.*

Proof. By Proposition 2.2(2), p-dimensional local Hopf algebras are in one to one correspondence with p-dimensional connected Hopf algebras by vector space dual. Therefore by Theorem 6.1, we know there are only two non-isomorphic p-dimensional local Hopf algebras. $\mathbf{k}[x]/(x^p)$ is local as p-dimensional algebra. With respect to the coalgebra structure, when $\Delta(x) = x \otimes 1 + 1 \otimes x$, it is connected. When $\Delta(x) = x \otimes 1 + 1 \otimes x + x \otimes x$, we have $\Delta(x+1) = (x+1) \otimes (x+1)$, which is a group-like element. Therefore it is cosemisimple. They are certainly non-isomorphic as coalgebras. $\square$

In the rest of the section we focus on the structure of connected Hopf algebras of dimension p^2. We first consider the case when $\dim \mathrm{P}(H) = 1$. By Remark 5.3, we have $\mathbf{k} \subset K \subset H$ where K is generated by some $x \in \mathrm{P}(H)$. Since $K \subsetneq H$, by Theorem B.6, there exists some $y \in H \setminus K$ such that $(\Delta(y) - y \otimes 1 - 1 \otimes y)$ is a nonzero cohomology class in $\mathrm{H}^2(\Omega K)$. Furthermore according to Lemma B.1, it is clear that $\dim \mathrm{H}^2(\Omega K) = 1$. Hence the nonzero cohomology class can be chosen in a unique way up to a scalar. It is easy to check that $\omega(x)=\sum_{i=1}^{p-1}\binom{p}{i}/p\ x^i\otimes x^{p-i}$ is a nonzero cohomology class in $\mathrm{H}^2(\Omega K)$.

Lemma 6.3. *Let H be a p^2-dimensional connected Hopf algebra with $dim\mathrm{P}(H)=1$. Then H is isomorphic to one the following*

1. $\mathbf{k}\left[x,y\right]/(x^p,y^p)$;
2. $\mathbf{k}\left[x,y\right]/(x^p,y^p-x)$;
3. $\mathbf{k}\left[x,y\right]/(x^p-x,y^p-y)$,

equipped with the comultiplications such that for x and y

$$\begin{aligned}\Delta(x) &= x\otimes 1+1\otimes x,\\ \Delta(y) &= y\otimes 1+1\otimes y+\omega(x).\end{aligned}\tag{6.0.3}$$

Proof. By the previous argument, we can find elements $x,y\in H$ with the stated comultiplications in (6.0.3). They generate a sub-bialgebra, hence a Hopf subalgebra by Remark 2.4, inside H. Since H has dimension p^2, H is generated by x,y. First we claim that x,y commute with each other. It is clear that $[x,y]$ is primitive since

$$\begin{aligned}\Delta\left([x,y]\right) &= [\Delta(x),\Delta(y)]\\ &= [x\otimes 1+1\otimes x, y\otimes 1+1\otimes y+\omega\left(x\right)]\\ &= [x,y]\otimes 1+1\otimes\left[x,y\right].\end{aligned}$$

In other words $[x,y]=\lambda x$ for some $\lambda \in \mathbf{k}$, which implies that $[x^n,y]=n\lambda x^n$ for any $n \geq 1$. Another calculation shows that

$$
\begin{aligned}
[\omega(x), y\otimes 1+1\otimes y] &= \left[\sum_{i=1}^{p-1}\binom{p}{i}/p\ x^i\otimes x^{p-i}, y\otimes 1+1\otimes y\right] \\
&= \sum_{i=1}^{p-1}\binom{p}{i}/p\ \left([x^i,y]\otimes x^{p-i}+x^i\otimes [x^{p-i},y]\right) \\
&= \sum_{i=1}^{p-1}\binom{p}{i}/p\ \left(i\lambda x^i\otimes x^{p-i}+x^i\otimes (p-i)\lambda x^{p-i}\right) \\
&= \sum_{i=1}^{p-1}\binom{p}{i}\lambda x^i\otimes x^{p-i} \\
&= 0.
\end{aligned} \tag{6.0.4}
$$

Since $\omega(x)^p=\omega(x^p)$ we have

$$\Delta\left(y^p\right)=\left(y\otimes 1+1\otimes y+\omega(x)\right)^p=y^p\otimes 1+1\otimes y^p+\omega(x^p). \tag{6.0.5}$$

By Theorem 6.1, we can assume that $x^p=0$ or $x^p=x$. When $x^p=0$, according to the above equation (6.0.5), y^p is primitive. Then we can write $y^p=\mu x$ for some $\mu\in\mathbf{k}$. Thus $\lambda^p x = x\mathrm{ad}(y)^p = [x,y^p]=[x,\mu x]=0$ which implies that $\lambda=0$. By further rescaling of the variables, μ can be taken either one or zero, which yields the first two non-isomorphic classes. On the other hand when $x^p=x$, we have y^p-y being primitive, where we can write $y^p=y+\mu x$ for some $\mu\in\mathbf{k}$. Therefore $[x,y]=[x^p,y]=\mathrm{ad}(x)^p y=0$. After the linear translation $y=y'+\sigma x$ satisfying $\sigma^p=\sigma+\mu$, we have $y'^p=y'$ while $\Delta(y')=y'\otimes 1+1\otimes y'+\omega(x)$. This completes the classification. □

Finally the classification for connected Hopf algebras of dimension p^2 follows:

Theorem 6.4. *Let H be a connected Hopf algebra of dimension p^2. When $\dim \mathrm{P}(H)=2$, it is isomorphic to one of the following:*

1. $\mathbf{k}\,[x,y]\,/\,(x^p,y^p)$;
2. $\mathbf{k}\,[x,y]\,/\,(x^p-x,y^p)$;
3. $\mathbf{k}\,[x,y]\,/\,(x^p-y,y^p)$;
4. $\mathbf{k}\,[x,y]\,/\,(x^p-x,y^p-y)$;
5. $\mathbf{k}\langle x,y\rangle/\left([xy]-y,x^p-x,y^p\right)$,

where x,y are primitive. When $\dim \mathrm{P}(H)=1$, it is isomorphic to one of the following:

(6) $\mathbf{k}\,[x,y]\,/(x^p,y^p)$;

(7) $\mathbf{k}\,[x,y]\,/(x^p,y^p-x)$;

(8) $\mathbf{k}\,[x,y]\,/(x^p-x,y^p-y)$,

where $\Delta\,(x)=x\otimes 1+1\otimes x$ *and* $\Delta\,(y)=y\otimes 1+1\otimes y+\omega(x)$.

Proof. By Proposition 2.2(7), when $\dim \mathrm{P}(H)=2$, it is primitively generated. Thus by Proposition 2.2(6), $H\cong u(\mathfrak{g})$ where $\dim\mathfrak{g}=2$. Therefore Proposition A.3 provides the classification for p^2-dimensional connected Hopf algebras in this case. And the other case when $\dim \mathrm{P}(H)=1$ directly comes from Lemma 6.3. □

Corollary 6.5. *Let* H *be a local Hopf algebra of dimension* p^2. *Then* H *is isomorphic to either* $\mathbf{k}\,[x,y]\,/\,(x^p,y^p)$ *or* $\mathbf{k}\,[x]\,/(x^{p^2})$ *as an algebra. Regarding the coalgebra structure for the first one we have the following distinct comultipications for* x,y:

(1) $\Delta\,(x)=x\otimes 1+1\otimes x$,
$\Delta\,(y)=y\otimes 1+1\otimes y$;

(2) $\Delta\,(x)=x\otimes 1+1\otimes x+x\otimes x$,
$\Delta\,(y)=y\otimes 1+1\otimes y$;

(3) $\Delta\,(x)=x\otimes 1+1\otimes x$,
$\Delta\,(y)=y\otimes 1+1\otimes y+\omega\,(x)$;

(4) $\Delta\,(x)=x\otimes 1+1\otimes x+x\otimes x$,
$\Delta\,(y)=y\otimes 1+1\otimes y+y\otimes y$;

(5) $\Delta\,(x)=x\otimes 1+1\otimes x+x\otimes x$,
$\Delta\,(y)=y\otimes 1+1\otimes y+x\otimes y$.

While for the second one we have

(6) $\Delta\,(x)=x\otimes 1+1\otimes x$;

(7) $\Delta\,(x)=x\otimes 1+1\otimes x+\omega\,(x^p)$;

(8) $\Delta\,(x)=x\otimes 1+1\otimes x+x\otimes x$.

Proof. By Proposition 2.2(2) and above Theorem 6.4, when $\dim \mathrm{P}(H^*)=2$ there are only five non-isomorphic classes for such H. Furthermore from Proposition 2.2(5), we know that $\dim J/J^2=\dim \mathrm{P}(H^*)=2$, where J is the Jacobson radical of H. Hence according to [12, Theorem 14.4], H is isomorphic to $\mathbf{k}[x,y]/(x^p,y^p)$ as an algebra since it is commutative. It is easy to check the above given coalgebra structures are non-isomorphic which completes the classification. The same argument applies to the other case. Theorem 6.4 shows that when $\dim \mathrm{P}(H^*) = 1$, there are three non-isomorphic classes. Since $\dim J/J^2 = \dim \mathrm{P}(H^*) = 1$, H is isomorphic to $\mathbf{k}[x]/(x^{p^2})$ as algebras. Because those three given above are definitely non-isomorphic as coalgebras, they complete the list. $\square$

Theorem 6.6. *Let H be a finite-dimensional connected Hopf algebra with* $\mathrm{P}(H) = 1$*. Then H is local if and only if the primitive elements are nilpotent.*

Proof. We only prove one direction. Suppose $\dim H = p^n$ and let $0 \neq x \in \mathrm{P}(H)$. We prove inductively on n such that when x is nilpotent then H is local. The first two cases when $n = 1, 2$ follow from Theorem 6.1 and 6.4. Assume now that $n \geq 3$. By Remark 5.3, we can filter H by an increasing series of normal Hopf subalgebras:

$$\mathbf{k} = K_0 \subset K_1 \subset K_2 \subset \cdots \subset K_n = H$$

where K_1 is generated by x and $\dim K_m = p^m$ for all $0 \leq m \leq n$. As in the proof of Lemma 5.1, there exists an injection $K_{n-1}/K_1^+K_{n-1} \hookrightarrow K_n/K_1^+K_n$. It induces an isomorphism on the primitive spaces since both of them have dimension one by Proposition 5.2(4). By induction K_{n-1} is local and so is $K_{n-1}/K_1^+K_{n-1}$. Therefore $K_n/K_1^+K_n$ is local since the primitive elements are nilpotent by the above injection. Thus there exits integers l, s such that $(\ker \varepsilon)^l \subseteq K_1^+K_n$ and $(K_1^+)^s = 0$, where $\ker \varepsilon$ is the augmentation ideal of H. Hence $(\ker \varepsilon)^{ls} \subseteq (K_1^+K_n)^s = (K_1^+)^s K_n = 0$. So H is local. $\square$

Theorem 6.7. *Let H be a finite-dimensional connected Hopf algebra with* $\mathrm{P}(H) = 1$*. Then the center of H contains* $\mathrm{P}(H)$*.*

Proof. Let $0 \neq x \in \mathrm{P}(H)$. Suppose $\dim H = p^n$. We show by inductive on n that $x \in Z(H)$, the center of H. It is trivial when $n = 1$. Assume that $n \geq 2$. By Remark 5.3, we can filter H by an increasing series of normal Hopf subalgebras:

$$\mathbf{k} = K_0 \subset K_1 \subset K_2 \subset \cdots \subset K_n = H$$

satisfying K_1 is generated by x and $K_{n-1} \subset H$ is a degree one normal extension. Then by Theorem B.6, we can find some $y \in H \setminus K_{n-1}$ such that $\Delta(y) = y \otimes 1 + 1 \otimes y + u$ where $u \in K_{n-1} \otimes K_{n-1}$, which together with K_{n-1} generate H. Moreover we have the relation $y^p + \lambda y + a = 0$ for some $\lambda \in \mathbf{k}$ and $a \in K_{n-1}$.

By induction $x \in Z(K_{n-1})$. Then it suffices to show $[x, y] = 0$. It is easy to check that $[x, y]$ is primitive. Therefore we can write $[x, y] = \mu x$ for some $\mu \in \mathbf{k}$. By Theorem 6.1, we can further assume either $x^p = 0$ or $x^p = x$. When $x^p = 0$, by Theorem 6.6, H is local. Then its quotient $H/K_{n-1}^{+}H$, which is generated by the image of y, is local too. Hence the image of y in $H/K_{n-1}^{+}H$ is nilpotent since it is primitive. Thus in the relation of y we must have $\lambda = 0$ and $y^p + a = 0$. A calculation therefore shows that $\mu^p x = x(\mathrm{ad}y)^p = [x, y^p] = [x, -a] = 0$ which implies that $[x, y] = \mu x = 0$. When $x^p = x$, we have $[x, y] = [x^p, y] = (\mathrm{ad}x)^p y = 0$. This completes the proof. $\square$

Appendix A. Restricted Lie Algebras

We state the following technical lemma which is the key to our classification of finite-dimensional connected Hopf algebras.

Lemma A.1. [6, P. 186-187] *For any associative algebra A we have*

$$(x + y)^p = x^p + y^p + \sum_{i=1}^{p-1} s_i(x, y)$$

where $is_i(x, y)$ is the coefficient of λ^{i-1} in $x\left(\mathrm{ad}(\lambda x + y)\right)^{p-1}$ and

$$[x^p, y] = \left(\mathrm{ad}(x)\right)^p (y)$$

for any $x, y \in A$.

Definition A.2. A Lie algebra $\mathfrak{g}$ over $\mathbf{k}$ is restricted if there exists a map $\mathfrak{g} \to \mathfrak{g}$, i.e., $x \mapsto x^{[p]}$ satisfying, for all $x, y \in \mathfrak{g}$ and $\alpha \in \mathbf{k}$:

1. $(\alpha x)^{[p]} = \alpha^p x^{[p]}$,
2. $(x + y)^{[p]} = x^{[p]} + y^{[p]} + \sum_{i=1}^{p-1} s_i(x, y)$, where $is_i(x, y)$ is the coefficient of λ^{i-1} in $x\left(\mathrm{ad}(\lambda x + y)\right)^{p-1}$,
3. $\left[x, y^{[p]}\right] = x\left(\mathrm{ad}(y)\right)^p$.

If $\mathfrak{g}$ is restricted and $\mathcal{U}(\mathfrak{g})$ is the usual enveloping algebra, let B be the ideal in $\mathcal{U}(\mathfrak{g})$ generated by all $x^p - x^{[p]}, x \in \mathfrak{g}$, and define $u(\mathfrak{g}) = \mathcal{U}(\mathfrak{g})/B$. Then $u(\mathfrak{g})$ is called the restricted enveloping algebra of $\mathfrak{g}$. A version of the PBW theorem holds for $u(\mathfrak{g})$: given a basis for $\mathfrak{g}$, the ordered monomials in this basis, where the exponent of each basis element is bounded by $p-1$, form a basis for $u(\mathfrak{g})$. Consequently if $\dim \mathfrak{g} = n$, then $\dim u(\mathfrak{g}) = p^n$.

Now denote $\mathfrak{g}$ as a two-dimensional Lie algebra with basis $\{x, y\}$. It is well known that there is only one nonabelian two-dimensional Lie algebra up to isomorphisms, where we can write $[x, y] = y$.

Proposition A.3. *Let $\mathfrak{g}$ be a two-dimensional restricted Lie algebra with basis $\{x, y\}$. Then the restricted maps can be classified as follows: When $\mathfrak{g}$ is abelian:*

(1) $x^{[p]} = 0, y^{[p]} = 0$;
(2) $x^{[p]} = x, y^{[p]} = 0$;
(3) $x^{[p]} = y, y^{[p]} = 0$;
(4) $x^{[p]} = x, y^{[p]} = y$.

When $\mathfrak{g}$ is nonabelian such that $[x, y] = y$:

(5) $x^{[p]} = x, y^{[p]} = 0$.

Proof. Suppose that $\mathfrak{g}$ is abelian. First we will prove that there exists a restricted Lie subalgebra of dimension one. Assume that the restricted map of $\mathfrak{g}$ is given by:

$$\begin{cases} x^{[p]} = a_{11}x + a_{12}y \\ y^{[p]} = a_{21}x + a_{22}y. \end{cases} \tag{A.0.6}$$

For any element in $\mathfrak{g}$, say $z = \lambda x + \mu y$ where $\lambda, \mu \in \mathbf{k}$ we have:

$$z^{[p]} = \lambda^p x^{[p]} + \mu^p y^{[p]} = \left(a_{11}\lambda^p + a_{21}\mu^p\right) x + \left(a_{12}\lambda^p + a_{22}\mu^p\right) y.$$

If $z^{[p]} = z$ then all the coefficients satisfy the following equations:

$$\begin{cases} a_{11}\lambda^p + a_{21}\mu^p = \lambda \\ a_{12}\lambda^p + a_{22}\mu^p = \mu. \end{cases}$$

When $a_{21} \neq 0$, we have $\mu^p = \left(\lambda - a_{11}\lambda^p\right)/a_{21}$ which implies the following equation:

$$\left(a_{12}a_{21} - a_{11}a_{22}\right)^p / a_{21}^p \lambda^{p^2} + \left(a_{11}/a_{21} + a_{22}^p/a_{21}^p\right)\lambda^p - 1/a_{21}\lambda = 0.$$

Furthermore if $a_{12}a_{21} - a_{11}a_{22} \neq 0$, then there exist solutions λ, μ not all zero such that $z^{[p]} = z$. Otherwise if $a_{12}a_{21} - a_{11}a_{22} = 0$ and $a_{21} \neq 0$, we can choose $\mu \neq 0$ such that $(x + \mu y)^{[p]} = 0$ where

$\mu^p = -a_{11}/a_{21}$. Finally if $a_{21} = 0$, set $z = y$ by rescaling we have $z^{[p]} = z$ or $z^{[p]} = 0$. Thus without loss of generality we can replace (A.0.6) by

$$\begin{cases} x^{[p]} = ax + by \\ y^{[p]} = cy. \end{cases}$$

If $a = c = 0$, then there are two cases either $b = 0$ or $b \neq 0$. By rescaling we have either $x^{[p]} = 0, y^{[p]} = 0$ or $x^{[p]} = y, y^{[p]} = 0$. If $a \neq 0, c \neq 0$, after taking a linear transformation

$$\begin{cases} x = dx' + ey' \\ y = fy', \end{cases}$$

where d, e, f satisfy the following equations whose solutions are easy to check:

$$\begin{cases} d^p = ad \\ e^p cf = f^p(ae + bf) \\ f^p = cf \end{cases}$$

would give case (4). The remaining possibilities when $a = 0, c \neq 0$ or $a \neq 0, c = 0$, by linear transformations, would give case (2). At last when $\mathfrak{g}$ is nonabelian, by the third condition of the definition of restricted Lie algebras, we have $[x, x^{[p]}] = [y, y^{[p]}] = [x, y^{[p]}] = 0$ and $[x^{[p]}, y] = y$. Since $[x, y] = y$, then $x^{[p]} = x, y^{[p]} = 0$. $\square$

Appendix B. Cohomology of Coalgebras

Let C be a coaugmented coalgebra in the sense that there is a coalgebra map from the trivial coalgebra $\mathbf{k}$ to C. Denote $J = C^+$, the kernel of the counit, and one defines the reduced comultiplication on J by

$$\overline{\Delta}(c) = \Delta(c) - (1 \otimes c + c \otimes 1). \tag{B.0.7}$$

Then the cobar construction ΩC on C is the differential graded algebra defined as follows:

- As a graded algebra, ΩC is the tensor algebra $T(J)$,
- The differential in ΩC is given by

$$\partial_C^n = \sum_{i=0}^{n-1} (-1)^i \mathrm{Id}^{\otimes i} \otimes \overline{\Delta} \otimes \mathrm{Id}^{\otimes (n-i-1)}. \tag{B.0.8}$$

Dually given an augmented algebra A, one can construct a differential graded coalgebra BA, which is called the bar construction of A. See [3, §19] for the basic properties of cobar and bar constructions.

Lemma B.1. *Let C be a finite-dimensional coalgebra and A be its dual algebra. Then* $\dim \mathrm{H}^n(\Omega C) = \dim \mathrm{Ext}^n_A(\mathbf{k}, \mathbf{k})$ *for every $n \geq 0$.*

Proof. When C is finite-dimensional, denote A as its dual algebra. According to [7, Lemma 8.6(c)], $B^{\#}A \cong \Omega C$ as DG algebras, where $B^{\#}A$ is the graded dual of the bar construction of A. Meanwhile according to [8, Lemma 4.2], $\mathrm{H}^{\bullet}(B^{\#}A) \cong \mathrm{Ext}^{\bullet}_A(\mathbf{k}, \mathbf{k})$. This gives the proof. $\square$

Proposition B.2. *Let $\mathfrak{g}$ be a restricted Lie algebra with basis $\{x_1, x_2, \cdots, x_n\}$ and denote $K = u(\mathfrak{g})$. Then $\{\omega(x_i),\ x_j \otimes x_k | 1 \leq i \leq n, 1 \leq j < k \leq n\}$ is a basis for $\mathrm{H}^2(\Omega K)$ with all x_i being primitive.*

Proof. By Lemma B.1, $\dim \mathrm{H}^2(\Omega K) = \dim \mathrm{Ext}^2_{K^*}(\mathbf{k}, \mathbf{k}) = n(n+1)/2$, since the dual K^* is isomorphic to $\mathbf{k}[y_1, \cdots, y_n]/(y_1^p, \cdots, y_n^p)$ as algebras. Notice all $\omega(x_i)$ and $\{x_j \otimes x_k | 1 \leq j < k \leq n\}$ are cocycles. We know K has a basis $\{x_1^{i_1} x_2^{i_2} \cdots x_n^{i_n} | 0 \leq i_1, i_2, \cdots, i_n \leq p-1\}$, it is easy to argue the linear independence. $\square$

Lemma B.3. *Let $\mathfrak{g}$ be a restricted Lie algebra and denote $K = u(\mathfrak{g})$. Then*

$$\left(\omega(\sum_{i=1}^{n} \lambda_i x_i) - \sum_{i=1}^{n} \lambda_i^p \omega\left(x_i\right)\right)$$

is a zero cohomology class in $\mathrm{H}^2(\Omega K)$, where $x_i \in \mathfrak{g}$ for all $1 \leq i \leq n$.

Proof. Without loss of generality we can assume $\mathfrak{g}$ is finite-dimensional. Because $\mathbf{k}$ is algebraically closed over $\mathbb{F}_p$, we can replace $\mathbf{k}$ with some finite field $\mathbb{F}_q$. By basic algebraic number theory, there exists some number field L, where p remains prime in the ring of integers $\mathcal{O}_L$ and $\mathcal{O}_L/p = \mathbb{F}_q$. Now by choosing the representatives for $\mathbb{F}_q$ in $\mathcal{O}_L$, we can consider $\mathfrak{g}$ as a free module over $\mathcal{O}_L$ with a bilinear bracket $[,]$ representing all the relations between the basis. Denote $A = U(\mathfrak{g})$ as the enveloping algebra over $\mathcal{O}_L$. There is a quotient map $\pi : A \twoheadrightarrow K$ which factors through $A/(p)$. Therefore it suffices to prove that

$$(\omega(x+y) - \omega(x) - \omega(y))$$

is a zero cohomology class in $\mathrm{H}^2(\Omega A)$ for any $x, y \in \mathfrak{g}$. Then the general result follows by using the quotient map π and induction. Apply Lemma A.1 in $A \otimes_{\mathcal{O}_L} \mathcal{O}_L/(p)$, we have $(x+y)^p \equiv x^p + y^p + z$

(mod p) for some $z \in \mathfrak{g}$. Therefore $(x+y)^p = x^p + y^p + z + p\Theta$ for some $\Theta \in A$ and

$$\begin{aligned}
\Delta(x+y)^p &= (\Delta(x+y))^p \\
&= ((x+y) \otimes 1 + 1 \otimes (x+y))^p \\
&= (x+y)^p \otimes 1 + 1 \otimes (x+y)^p + p\omega(x+y).
\end{aligned}$$

On the other hand

$$\begin{aligned}
\Delta(x+y)^p &= \Delta(x^p + y^p + z + p\Theta) \\
&= x^p \otimes 1 + 1 \otimes x^p + p\omega(x) + y^p \otimes 1 + 1 \otimes y^p + p\omega(y) + z \otimes 1 + 1 \otimes z + p\Delta(\Theta) \\
&= (x^p + y^p + z) \otimes 1 + 1 \otimes (x^p + y^p + z) + p\omega(x) + p\omega(y) + p\Delta(\Theta).
\end{aligned}$$

Thus $p\partial_A^1(\Theta) = p\left\{\omega(x+y) - \omega(x) - \omega(y)\right\}$. Finally $\partial_A^1(\Theta) = \omega(x+y) - \omega(x) - \omega(y)$. $\square$

Definition B.4. Let H be a Hopf algebra. Define the degree zero adjoint map T_x on ΩH by

$$T_x^n = \sum_{i=0}^{n-1} \mathrm{Id}^{\otimes i} \otimes \mathrm{ad}(x) \otimes \mathrm{Id}^{\otimes(n-i-1)}. \tag{B.0.9}$$

where $\mathrm{ad}(x)(H) = [x, H]$ for any $x \in H$.

Lemma B.5. *For every Hopf algebra H, T_x defines a chain map from ΩH to itself for all $x \in \mathrm{P}(H)$. Furthermore $H^\bullet(\Omega H)$ becomes a graded $\mathfrak{g}$-module through the adjoint map where $\mathfrak{g} = \mathrm{H}^1(\Omega H)$.*

Proof. We prove inductively that $\partial^n T^n = T^{n+1}\partial^n$ for all $n \geq 0$. It is clear when $n = 0, 1$ since $\overline{\Delta}\mathrm{ad}(x) = (\mathrm{ad}(x) \otimes 1 + 1 \otimes \mathrm{ad}(x))\,\overline{\Delta}$ for all $x \in \mathrm{P}(H)$. Suppose it is true for $n-1$ and then:

$$\begin{aligned}
\partial^n T^n &= \left(\partial^{n-1} \otimes \mathrm{Id} + (-1)^{n-1}\mathrm{Id}^{\otimes(n-1)} \otimes \partial^1\right)\left(T^{n-1} \otimes \mathrm{Id} + \mathrm{Id}^{\otimes(n-1)} \otimes T^1\right) \\
&= \partial^{n-1}T^{n-1} \otimes \mathrm{Id} + \partial^{n-1} \otimes T^1 + (-1)^{n-1}T^{n-1} \otimes \partial^1 + (-1)^{n-1}\mathrm{Id}^{\otimes(n-1)} \otimes \partial^1 T^1 \\
&= T^n\partial^{n-1} \otimes \mathrm{Id} + \partial^{n-1} \otimes T^1 + (-1)^{n-1}T^{n-1} \otimes \partial^1 + (-1)^{n-1}\mathrm{Id}^{\otimes(n-1)} \otimes T^2\partial^1 \\
&= T^n\partial^{n-1} \otimes \mathrm{Id} + \partial^{n-1} \otimes T^1 + (-1)^{n-1}\left(T^{n-1} \otimes \mathrm{Id}^{\otimes 2} + \mathrm{Id}^{\otimes(n-1)} \otimes T^1 \otimes \mathrm{Id}\right)\left(\mathrm{Id}^{\otimes(n-1)} \otimes \partial^1\right) \\
&\quad + (-1)^{n-1}\mathrm{Id}^{\otimes(n-1)} \otimes (\mathrm{Id} \otimes T^1)\partial^1 \\
&= T^n\partial^{n-1} \otimes \mathrm{Id} + \partial^{n-1} \otimes T^1 + (-1)^{n-1}\left(T^n \otimes \mathrm{Id}\right)\left(\mathrm{Id}^{\otimes(n-1)} \otimes \partial^1\right) + (-1)^{n-1}\mathrm{Id}^{\otimes(n-1)} \otimes \left(\mathrm{Id} \otimes T^1\right)\partial^1 \\
&= \left(T^n \otimes \mathrm{Id} + \mathrm{Id}^{\otimes n} \otimes T^1\right)\left(\partial^{n-1} \otimes \mathrm{Id} + (-1)^{n-1}\mathrm{Id}^{\otimes(n-1)} \otimes \partial^1\right) \\
&= T^{n+1}\partial^n
\end{aligned}$$

Then T induces an action of $\mathfrak{g} = \mathrm{P}(H)$ on $H^\bullet(\Omega H)$. A straightforward computation shows that $T_xT_y - T_yT_x = T_{[xy]}$ for any $x, y \in \mathrm{P}(H)$, which makes $H^\bullet(\Omega H)$ into a graded $\mathfrak{g}$-module. Finally $\mathrm{P}(H) \cong \mathrm{H}^1(\Omega H)$ by definition. □

Theorem B.6. *Let $K \subseteq H$ be an extension of connected Hopf algebras with first order $n \geq 2$. Then the differential ∂_H^1 induces an injective $\mathfrak{g}$-module map from H_n/K_n to $\mathrm{H}^2(\Omega K)$ where $\mathfrak{g} = \mathrm{P}(H)$.*

Proof. First of all by Corollary 3.3, H_n becomes a $\mathfrak{g}$-module by adjunction since $[H_1, H_n] \subseteq H_n$. Then choose $x \in H_n$ and we have $\Delta(x) = x \otimes 1 + 1 \otimes x + u$, where $u \in H_{n-1} \otimes H_{n-1} = K_{n-1} \otimes K_{n-1} \subseteq K \otimes K$ by [10, Lemma 5.3.2]. We know $(\Omega K, \partial_K)$ can be viewed as a sub-complex of $(\Omega H, \partial_H)$. Hence $\partial_K^2(u) = \partial_K^2\partial_H^1(x) = \partial_H^2\partial_H^1(x) = 0$, i.e., u is a cocycle. Then ∂_H^1 defines a linear map from H_n to $\mathrm{H}^2(\Omega K)$, which factors through H_n/K_n since $\partial_K^2\partial_K^1(K_n) = 0$. It is compatible with the $\mathfrak{g}$ action on $\mathrm{H}^2(\Omega K)$ by Lemma B.5. At last it is an injective map since if $\partial_H^2(x)$ were a coboundary for some $x \in H_n$, there would exist some $y \in K$ such that $\partial_K^1(y) = \Delta(y) - y \otimes 1 - 1 \otimes y = \partial_H^1(x)$. Hence $x - y \in H_1 = K_1$, which implies that $x \in K \bigcap H_n = K_n$ by Remark 2.4. □

Proposition B.7. *The first order for the extension from a restricted enveloping algebra can be $1, 2, p$ or ∞.*

Proof. Let $K = u(\mathfrak{g}) \subseteq H$ be one of such extensions, where $\mathfrak{g}$ is some restricted Lie algebra. The first order is one implies that $\mathfrak{g} \subsetneq \mathrm{P}(H)$ and if it is infinity means that $u(\mathfrak{g}) = H$. In the other cases we can assume that $K \neq H$ and the first order n is greater or equal to two. Denote $\{x_i\}$ as a basis for $\mathfrak{g}$. Then by Lemma B.6 and Proposition B.2, we can find some $x \in H_n \setminus K_n$ such that $\Delta(x) = x \otimes 1 + 1 \otimes x + \sum \alpha_i \omega(x_i) + \sum_{i<j} \alpha_{ij} x_i \otimes x_j$ where the coefficients are not all zero. Finally because the associated graded ring of H with respected to its coradical filtration is coradically graded as mentioned in [2, Def. 1.13], it is see to see that if all $\alpha_i = 0$ then $n = 2$. Otherwise $n = p$. □

References

[1] N. Andruskiewitsch and H.J. Schneider. Finite quantum groups and cartan matrices. *Advances in Mathematics*, 154(1):1–45, 2000.

[2] N. Andruskiewitsch and H.J. Schneider. Pointed hopf algebras. In *In New directions in Hopf algebras, MSRI series Cambridge Univ*, pages 1–68. Press, 2002.

[3] Y. Félix, S. Halperin, and J.C. Thomas. *Rational Homotopy Theory*. Number 110 in Graduate Texts in Mathematics. Springer-Verlag, New York, second edition, 2001.

[4] J.E. Humphreys. *Introduction to Lie algebras and representation theory*, volume 9. Springer, 2000.

[5] I.M. Isaacs. *Algebra: a graduate course*, volume 100. Amer Mathematical Society, 1994.

[6] N. Jacobson. *Lie Algebras*. Dover Publications, Inc., New York, 1979.

[7] D.-M. Lu, J.H. Palmieri, Q.-S. Wu, and J.J. Zhang. A-infinity algebras for ring theorists. *Algebra Colloquium*, 11(1), 2004.

[8] D.-M. Lu, J.H. Palmieri, Q.-S. Wu, and J.J. Zhang. A-infinity structure on ext-algebras. *Journal of Pure and Applied Algebra*, 213:2017–2037, 2009.

[9] J.W. Milnor and J.C. Moore. On the structure of hopf algebras. *The Annals of Mathematics*, 81:211–264, Mar. 1965.

[10] S. Montgomery. *Hopf Algebras and Their Actions on Rings*. Number 82 in CBMS Regional Conference Series in Mathematics. American Mathematical Society, Providence, Rhode Island, 1993.

[11] D.G. Wang, JJ Zhang, and G. Zhuang. Lower bounds of growth of hopf algebras. *Arxiv preprint arXiv:1101.1116*, 2011.

[12] W.C. Waterhouse. *Introduction to Affine Group Schemes*. Number 66 in Graduate Texts in Mathematics. Springer-Berlag, New York, 1979.

[13] G. Zhuang. On the gelfand-kirillov dimension of pointed hopf algebras. inprint.